

## The effect of platelet-rich plasma on cavernous nerve regeneration in a rat model

Xie-Gang Ding,<sup>1</sup> Shi-Wen Li,<sup>1,\*</sup> Xin-Min Zheng,<sup>1</sup> Li-Quan Hu,<sup>1</sup> Wan-Li Hu,<sup>1</sup> and Yi Luo<sup>1</sup>

<sup>1</sup>Research Center of Urology and Andrology, Zhongnan Hospital, Wuhan University, Wuhan 430071, China

\*Research Center of Urology and Andrology, Zhongnan Hospital, Wuhan University, Wuhan 430071, China.

Fax: +86-27-6781-3090 E-mail: [drlishiwen@yahoo.com.cn](mailto:drlishiwen@yahoo.com.cn)

Received 2008 Jul 25; Revised 2008 Oct 6; Accepted 2008 Oct 22.

Copyright © 2009 SIMM & SJTU

This article has been [cited by](#) other articles in PMC.

### Abstract

[Go to:](#)

The aim of this study was to investigate the effect of platelet-rich plasma (PRP) on cavernous nerve (CN) regeneration and functional status in a nerve-crush rat model. Twenty-four Sprague–Dawley male rats were randomly divided into three equal groups: eight had a sham operation, eight underwent bilateral nerve crushing with no further intervention and eight underwent bilateral nerve crushing with an immediate application of PRP on the site of injury. Erectile function was assessed by CN electrostimulation at 3 months and nerve regeneration was assessed by toluidine blue staining of CN and nicotinamide adenine dinucleotide phosphate (NADPH)-diaphorase staining of penile tissue. Three months after surgery, in the group that underwent bilateral nerve crushing with no further intervention, the functional evaluation showed a lower mean maximal intracavernous pressure (ICP) and maximal ICP per mean arterial pressure (MAP) with CN stimulation than those in the sham group. In the group with an immediate application of PRP, the mean maximal ICP and maximal ICP/MAP were significantly higher than those in the injured control group. Histologically, the group with the application of PRP had more myelinated axons of CNs and more NADPH-diaphorase-positive nerve fibres than the injured control group but fewer than the sham group. These results show that the application of PRP to the site of CN-crush injury facilitates nerve regeneration and recovery of erectile function. Our research indicates that clinical application of PRP has potential repairing effect on CN and peripheral nerves.

**Keywords:** platelet rich plasma, impotence, erectile dysfunction, nerve regeneration

### Introduction

[Go to:](#)

With the development of the anatomic radical prostatectomy (RP) described by Walsh and Donker<sup>1</sup>, impotence rates following RP have decreased significantly. However, even now, a significant proportion of the patients undergoing anatomic nerve-sparing RP often experience incomplete or delayed recovery of erectile function.

In many instances, the cavernous nerves (CNs) may have been inadvertently damaged by manipulation during nerve-sparing prostatectomy. The recovery of erectile function may depend on re-growth of nerves from the remaining neural tissue<sup>2</sup>. Accumulating evidence indicates that neuroimmunophilin ligand and many growth factors, such as insulin growth factor-1 (IGF-1), brain-derived growth factor (BDNF) and vascular endothelial growth factor (VEGF) play a significant role in neural regeneration and up-regulation of neuronal nitric oxide synthase (nNOS), as well as in the recovery of erectile function after CN injury<sup>3, 4, 5, 6, 7, 8, 9, 10</sup>.

Platelet-rich plasma (PRP) is prepared by centrifugation of the patient's own blood <sup>11</sup>. Platelets contain various growth factors, such as platelet-derived growth factor (PDGF), transforming growth factor (TGF)- $\beta$ , IGF-I and VEGF. When platelets are activated, they release those factors, which play important biological roles in various conditions <sup>12</sup>. These findings suggest that the application of PRP to the injured CN could facilitate nerve repair.

The aim of this study was to evaluate whether the application of PRP to the site of CN injury had neuromodulatory effects on the regeneration of CNs and also to evaluate the recovery of erectile function after bilateral CN-crush injury in a rat model.

## Materials and methods

[Go to:](#)

### Experimental animals

Twenty-four healthy male Sprague-Dawley rats (3 months old, 250–300 g) were randomly divided into three equal groups: group 1 received a sham operation comprising a midline incision and identification of the CNs bilaterally, with no further surgical manipulation; group 2 underwent CN dissection and a subsequent intentional 2-min crush injury with a hemostatic clamp; and group 3 was treated as group 2, but followed by an immediate application of activated PRP to the site of CN injury. All rats were obtained from the Experimental Animal Centre of Wuhan University (Wuhan, China), and the Wuhan University Animal Care and Use Committee approved all the procedures followed in this study.

### PRP preparation

The PRP preparation was obtained from six age-matched male Sprague-Dawley rats. The whole blood of six rats was drawn preoperatively through cardiac puncture into ethylene diamine tetraacetic acid-coated tubes. The blood was subjected to centrifugation for 10 min at  $312 \times g$  and the supernatant was transferred to another tube. The supernatant was subjected to centrifugation for 10 min at  $1\,248 \times g$  to yield platelet-poor plasma (PPP) and PRP. The top layer, which consisted of the PPP, was aspirated and put into a new tube. The remaining layer was the PRP. Approximately 400  $\mu\text{L}$  of PRP was activated with 40  $\mu\text{L}$  of 10% calcium chloride solution and 1 000 U  $\text{mL}^{-1}$  bovine thrombin (Sigma-Aldrich, St. Louis, MO, USA) to form a gel for application to every rat in group 3 during the operation.

### Surgical procedures

For the surgical procedure, rats were anaesthetized by an intraperitoneal injection with sodium pentobarbital (40 mg  $\text{kg}^{-1}$ ). The rats were kept isothermic by placing them on a heating pad at 37°C. A lower midline abdominal incision was made after the abdomen was shaved and prepared with an iodine-based solution. With the aid of an operating microscope (SXP-1C, Medical Optical Instruments Factory of Shanghai Medical Instruments Co. Ltd, Shanghai, China), the prostate gland was exposed and the posterolateral CNs and major pelvic ganglion were identified bilaterally. There was no further surgical manipulation in group 1. In group 2, the CNs were isolated and a crush injury was applied using a hemostat clamp (HC-X020, Cheng-He Microsurgical Instruments Factory, Ningbo, China) for 2 min, and then the abdomen was closed. In group 3, after bilateral nerve crushing, 200  $\mu\text{L}$  of activated PRP was applied immediately to each site of the CN injury. In all rats, the abdomen was closed in two layers.

### Measurement of erectile responses

The erectile response was assessed in all rats after 3 months by electrostimulation of the CNs and by measuring intracavernous pressure (ICP). Through a repeat midline abdominal incision, the CNs were exposed and isolated. The skin overlying the penis was incised and the crura of the penis was identified. A 23-G scalp-vein needle filled with 250 U  $\text{mL}^{-1}$  of heparin solution was connected to polyethylene-50 tubing and inserted into the right crus body to measure the ICP. Systemic mean arterial pressure (MAP) was monitored by inserting a 22-G

cannula into the carotid artery on the left side of the incised neck. A bipolar stainless steel electrode was used to directly stimulate the CNs (probes 2 mm in diameter and separated by 1 mm). Monophasic rectangular pulses were generated by a computer with a custom-built constant current amplifier. The stimulus parameters were as follows: amplitude, 1.5 mA; frequency, 20 Hz; pulse width, 0.2 ms; and duration, 50 s. The ICPs and MAP were recorded in all rats using a bioinformation acquisition system (BL-420F, Chengdu TME Technology Co. Ltd, Chengdu, China). The mean of the maximum right and left ICPs was then determined for each rat. Finally, the maximal ICP/MAP ratios in the three groups were calculated.

After functional testing, samples of CNs and the mid-shaft penis were collected for toluidine blue staining and nicotinamide adenine dinucleotide phosphate (NADPH)-diaphorase staining.

### **Toluidine blue staining**

Two to three mm of CN was obtained from the main trunk of both sides at equivalent points, which were distal to the site of injury. All harvested nerves were fixed in 3% (weight/volume) cold glutaraldehyde. The nerve segments were dehydrated with ethanol and postfixed with 1% osmium tetroxide before being infiltrated with a graded araldite-propylene oxide mixture and embedded in EPON 812 (Heidelberg, NY, USA). An LKB III Ultramicrotome (LKB Produkter A.B., Broma, Sweden) was used to obtain 1  $\mu\text{m}$  cross-sections of the embedded nerve, which were then stained with 1% toluidine blue and examined with light microscopy. Images were captured using an Olympus-DP12 camera ( $\times 100$  oil immersion objective) and processed using Image Pro-Plus 3.0 (MediaCybernetics, Bethesda, MD, USA). The nerves were analyzed to determine the number of myelinated axons.

### **Nicotinamide adenine dinucleotide phosphate-diaphorase staining**

Samples of penile tissue were fixed for 4 h in phosphate buffer containing 0.002% picric acid and 2% formaldehyde, and then transferred to 30% sucrose before freezing. Serial cryosections (10  $\mu\text{m}$ ) were adhered to the charged slides. After being air dried for 5 min, the sections were incubated with 0.1  $\text{mmol L}^{-1}$  NADPH, 0.2  $\text{mmol L}^{-1}$  nitroblue tetrazolium and 0.2% Triton X-100 (Sigma-Aldrich) in a buffer, with constant microscopic monitoring for color development. When the medium was deep blue for NADPH-diaphorase-positive nerves, rinsing in buffer terminated the reaction. The staining pattern was assessed by counting the number of NADPH-diaphorase-positive nerves in the dorsal nerves at an original magnification of  $\times 400$  with light microscopy. The endothelium staining was excluded from the count.

### **Statistical analysis**

Data were expressed as mean  $\pm$  SD. The results were first analyzed using one-way analysis of variance (ANOVA) with significance indicated at  $P < 0.05$ . If the difference was significant, a Student–Newman–Keuls test was carried out.


## **Results**

[Go to:](#)


### **Evaluation of erectile function**

To evaluate recovery of erectile function, the maximal ICP was measured ([Figure 1](#)). No erectile dysfunction was identified in the sham group, with a pressure of  $108.9 \pm 12.8$   $\text{cmH}_2\text{O}$ . The injured control group showed a significant reduction in maximal ICP of  $32.7 \pm 12.2$   $\text{cmH}_2\text{O}$ , consistent with a state of erectile dysfunction. In the PRP-treated group, the mean maximal ICP was  $83.1 \pm 12.9$   $\text{cmH}_2\text{O}$ , which was significantly higher than that in the injured control group ( $P < 0.05$ ). The maximal ICP/MAP ratio was increased significantly in the PRP treated group compared with the injured control group, but it was still less than that of the sham group ([Figure 2](#)).

[Figure 1](#)



Example of Maximal intracavernous pressure (ICP) changes after electrostimulation of cavernous nerves at 3 months. (A): Group 1; the pressure reached 100 cmH<sub>2</sub>O. (B): Group 2; the pressure reached 38 cmH<sub>2</sub>O. (C): Group 3; the pressure reached 92 cmH<sub>2</sub>O.




**Figure 2**

Maximal intracavernous pressure (ICP)/mean arterial pressure (MAP) ratio with ICP responses by electrical stimulation in three groups. <sup>b</sup> $P < 0.05$ , compared with group 2; <sup>c</sup> $P < 0.05$ , compared with group 1.

### Toluidine blue staining of cavernous nerve

The myelinated axons of CNs were assessed by toluidine blue staining of histological specimens. In concordance with the functional result, the number of myelinated axons in the injured control group was significantly less than that in the sham group ( $P < 0.05$ ) (Table 1 and Figure 3). The group treated with PRP showed a statistically significant increase in the number of myelinated axons compared with the injured control group ( $P < 0.05$ ), but it was still less than that in the sham group ( $P < 0.05$ ).



**Figure 3**

Toluidine blue staining of cavernous nerves: (A): Group 1; (B): Group 2; (C): Group 3. In group 1, there were abundant myelinated axons of cavernous nerves that had a normal morphological appearance. In group 2, the number of myelinated axons decreased ...

Group	Myelinated axons (mean ± SD)	NADPH-diaphorase-positive nerve fibres (mean ± SD)
Sham	100 ± 10	100 ± 10
Injured control	30 ± 5	30 ± 5
PRP-treated	60 ± 10	60 ± 10

<sup>a</sup>  $P < 0.05$  compared with group 1.  
<sup>b</sup>  $P < 0.05$  compared with group 2.

**Table 1**


Changes in the number of myelinated axons of the CN and NADPH-diaphorase-positive nerve fibres in the dorsal nerve in the three groups of rats (mean ± SD).

### Nicotinamide adenine dinucleotide phosphate-diaphorase staining

There were significantly fewer NADPH-diaphorase-positive nerve fibres in the dorsal nerves of the injured control group than in the sham group ( $P < 0.05$ ) (Table 1 and Figure 4). In the PRP-treated group, the number of NADPH-diaphorase-positive nerve fibres was increased significantly compared with the injured control group ( $P < 0.05$ ).

**Figure 4**

NADPH-diaphorase staining of dorsal nerves. (A): Group 1; (B): Group 2; (C): Group 3. In group 1, there was predominant blue staining of nerve fibres. In group 2, there was a paucity of blue-stained fibres. In group 3, there was a significant increase ...



## Discussion

Go to:

Our results show a measurable neurotrophic effect of PRP on CN regeneration after crush injury in a rat model. This is on the basis of the following statistically significant results found when the injured CN was treated with PRP: (1) during CN electrostimulation, the measurement of the maximum ICP and ICP/MAP in this group showed good functional recovery after treatment; (2) histologic evaluation of this group showed a significant increase in the number of myelinated axons, which indicated the recovery of CNs directly; and (3) an increase of NADPH-diaphorase-positive nerve fibres in the dorsal nerves also showed the recovery of the CNs indirectly after application of PRP. These findings are in agreement with an earlier report of the neurotrophic and neuroprotective effects of PRP on facial nerve regeneration in a rat model [13](#).

The specific mechanism for regeneration of damaged CNs is still unknown, but growth factors are likely to play an important role in this process. Jung *et al.* [14](#) created a rat model of CN neurotomy to study changes in nNOS-containing nerve fibres and the RNA expression of IGF-I, nerve growth factor, TGF- $\alpha$ , TGF- $\beta_1$ ,  $\beta_2$  and  $\beta_3$  and NOS on the penis. Their results suggested that IGF-I and TGF- $\beta_2$  may contribute to the regeneration of nNOS-containing fibres. In a more recent study, Hu *et al.* [15](#) found that autologous vein grafts combined with local injection of IGF-I facilitated repair of ablated CNs. Researchers also found that intracavernous injection with VEGF and BDNF or adeno-associated virus-mediated BDNF enhanced nerve regeneration and erectile recovery after bilateral CN injury [8, 9, 10](#). In addition to *in vivo* studies in rats undergoing injury to CNs, Lin *et al.* [16](#) established an *in vitro* model of pelvic ganglion culture. Among several growth factors tested, they found that both BDNF and VEGF facilitated the outgrowth of axons that stained positive for nNOS. Other studies indicated that BDNF promoted neurite growth in the main pelvic ganglion, primarily by activating the Janus kinase–signal transducer and activator of the transcription (JAK–STAT) pathway [17, 18](#). Both the *in vivo* and the *in vitro* studies mentioned above showed the beneficial role of growth factors in neural regeneration and up-regulation of nNOS, as well as in the recovery of erectile function.

In this study, the application of PRP to the site of CN injury shows the positive effect on CN regeneration and functional recovery. We believe that the growth factors and platelet gel in PRP are all responsible for this property. The platelets in the PRP are activated by the addition of thrombin and excess calcium, which promotes both platelet activation and the formation of a thrombus-like gelatinous substance (platelet gel). The activated platelets are trapped on the platelet gel, where they continue to excrete growth factors and slowly diffuse into the injury site. Such a platelet gel could be a good delivery vehicle for the growth factors PDGF, TGF- $\beta$ , IGF-I and VEGF, which are present in PRP. The above-mentioned researchers have found that TGF- $\beta_2$ , IGF-I and VEGF contribute to the neural regeneration and up-regulation of nNOS. As for PDGF, it has been shown that neurons express PDGF receptors, and PDGF- $\beta$  has been proven to be a mitogen and survival factor for Schwann cells with trophic activity on neurons [19](#). Also, an augmented PDGF- $\beta$  expression in peripheral neurons has been found after peripheral nerve injury, suggesting a role for PDGF- $\beta$  in peripheral nerve regeneration [20](#). Another study found that PDGF improved regeneration and remyelination of the nervous system [21](#). In our study, we also found that the group treated with PRP showed a statistically significant increase in the number of myelinated axons compared with the injured control group. Maybe it is partly contributed to the PDGF in PRP. The growth

factors in PRP not only react individually but may also have a compound effect. Therefore, we believe that the neuromodulatory effect of PRP may be associated with local releasing growth factors at the injured site and promotion of the myelinated axon's regeneration.

In summary, this study shows that the application of PRP to the site of CN injury has neuromodulatory effects on the regeneration of CNs and the recovery of erectile function. Given the limitations of the study, future work will include finding the optimal dose of PRP and identifying the primary molecular signaling pathway in the process of nerve repair. Although much is yet to be determined, our study indicates the potential of a new clinical application for PRP in CN or other peripheral nerve regeneration.

## Acknowledgments

[Go to:](#)

This project was supported by the National Natural Science Foundation of China (No. 30600616).

## Footnotes

[Go to:](#)

Edited by Dr Ching-Shwun Lin

## References

[Go to:](#)

1. Walsh PC, Donker PJ. Impotence following radical prostatectomy: insight into etiology and prevention. *J Urol.* 1982;128:492–7. [[PubMed](#)]
2. Carrier S, Zvara P, Nunes L, Kour NW, Rehman J, et al. Regeneration of nitric oxide synthase-containing nerves after cavernous nerve neurotomy in the rat. *J Urol.* 1995;153:1722–7. [[PubMed](#)]
3. Valentine H, Chen Y, Guo H, McCormick J, Wu Y, et al. Neuroimmunophilin ligands protect cavernous nerves after crush injury in the rat: new experimental paradigms. *Eur Urol.* 2007;51:1724–31. [[PMC free article](#)] [[PubMed](#)]
4. Lagoda G, Jin L, Lehrfeld TJ, Liu T, Burnett AL. FK506 and sildenafil promote erectile function recovery after cavernous nerve injury through antioxidative mechanisms. *J Sex Med.* 2007;4:908–16. [[PubMed](#)]
5. McCullough AR. Rehabilitation of erectile function following radical prostatectomy. *Asian J Androl.* 2008;10:61–74. [[PubMed](#)]
6. Jung GW, Kwak JY, Yoon S, Yoon JH, Lue TF. IGF-I and TGF-beta2 have a key role on regeneration of nitric oxide synthase (NOS)-containing nerves after cavernous neurotomy in rats. *Int J Impot Res.* 1999;11:247–59. [[PubMed](#)]
7. Bochinski D, Hsieh PS, Nunes L, Lin GT, Lin CS, et al. Effect of insulin-like growth factor-1 and insulin-like growth factor binding protein-3 complex in cavernous nerve cryoablation. *Int J Impot Res.* 2004;16:418–23. [[PubMed](#)]
8. Lin CS, Ho HC, Chen KC, Lin G, Nunes L. *et al.* Intracavernosal injection of vascular endothelial growth factor induces nitric oxide synthase isoforms. *BJU Int.* 2002;89:955–60. [[PubMed](#)]
9. Hsieh PS, Bochinski DJ, Lin GT, Nunes L, Lin CS, et al. The effect of vascular endothelial growth factor and brain-derived neurotrophic factor on cavernosal nerve regeneration in a nerve-crush rat model. *BJU Int.* 2003;92:470–5. [[PubMed](#)]
10. Chen KC, Minor TX, Rahman NU, Ho HC, Nunes L, et al. The additive erectile recovery effect of brain-derived neurotrophic factor combined with vascular endothelial growth factor in a rat model of neurogenic impotence. *BJU Int.* 2005;95:1077–80. [[PubMed](#)]
11. Sumida E, Iwasaki Y, Akiyoshi K, Kasugai S. Platelet separation from whole blood in an aqueous two-phase system with water-soluble polymers. *J Pharmacol Sci.* 2006;101:91–7. [[PubMed](#)]
12. Garg AK, Gargene D, Peace I. Using platelet-rich plasma to develop an autologous membrane for growth factor delivery in dental implant therapy. *Dent Implantol Update.* 2000;11:41–4. [[PubMed](#)]
13. Farrag TY, Lehar M, Verhaegen P, Carson KA, Byrne PJ. Effect of platelet rich plasma and fibrin sealant on facial nerve regeneration in a rat model. *Laryngoscope.* 2007;117:157–65. [[PubMed](#)]

14. Jung GW, Kwak JY, Kim IH, Koo MY, Park JI, et al. The role of growth factor on regeneration of nitric oxide synthase (NOS)-containing nerves after cavernous neurotomy in the rats. *Int J Impot Res.* 1999;11:227–35. [[PubMed](#)]
15. Hu WL, Hu LQ, Li SW, Zheng XM. Reconstruction of erectile reflex circuit by autologous vein graft combined with use of insulin-like growth factor. *Zhonghua Yi Xue Za Zhi.* 2004;84:1280–2. [[PubMed](#)]
16. Lin G, Chen KC, Hsieh PS, Yeh CH, Lue TF, et al. Neurotrophic effects of vascular endothelial growth factor and neurotrophins on cultured major pelvic ganglia. *BJU Int.* 2003;92:631–5. [[PubMed](#)]
17. Bella AJ, Lin G, Tantiwongse K, Garcia M, Lin CS, et al. Brain-derived neurotrophic factor (BDNF) acts primarily via the JAK/STAT pathway to promote neurite growth in the major pelvic ganglion of the rat: part I. *J Sex Med.* 2006;3:815–20. [[PubMed](#)]
18. Lin G, Bella AJ, Lue TF, Lin CS. Brain-derived neurotrophic factor (BDNF) acts primarily via the JAK/STAT pathway to promote neurite growth in the major pelvic ganglion of the rat: part 2 *J Sex Med* 2006;3:821–7. discussion 828–9.9 [[PubMed](#)]
19. Eccleston PA, Funa K, Heldin CH. Expression of platelet-derived growth factor (PDGF) and PDGF alpha- and beta-receptors in the peripheral nervous system: an analysis of sciatic nerve and dorsal root ganglia. *Dev Biol.* 1993;155:459–70. [[PubMed](#)]
20. Oya T, Zhao YL, Takagawa K, Kawaguchi M, Shirakawa K, et al. Platelet-derived growth factor-b expression induced after rat peripheral nerve injuries. *Glia.* 2002;38:303–12. [[PubMed](#)]
21. Allamargot C, Pouplard-Barthelaix A, Fressinaud C. A single intracerebral microinjection of platelet-derived growth factor (PDGF) accelerates the rate of remyelination *in vivo*. *Brain Res.* 2001;918:28–39. [[PubMed](#)]